PATHOLOGY TESTS EXPLAINED Information about pathology tests to help everyone take control of their health and make the right decisions about their care.

WHAT YOU SHOULD KNOW ABOUT SCREENING FOR CERVICAL CANCER

Cervical cancer is caused by the human papilloma virus – HPV. The virus infection is very common in both men and women, especially in people who are sexually active. Although it causes few symptoms and usually clears up naturally — just as the body recovers from cold and flu viruses – a few of the more than 100 types of the HP virus have the potential to cause cancer. HPV infection can lead to cancer of the cervix as well as oral, anal, and penile cancers.

The HPV test is used to detect these high-risk strains of HPV and if it shows that you have one of them, your sample will go on to be examined under the microscope to look for pre-cancerous changes or cancer. The results of this test will decide whether you need to be monitored or actively investigated.



l

What to expect when you have HPV testing

A sample of cells is taken from the cervical area during a pelvic examination, using a type of spatula, swab, or brush. The cells are transferred to a special preservative liquid and are transported to the laboratory for HPV testing.

The high level of accuracy of the test means that women between the ages of 25-75 only need to repeat the HPV test every five years after their initial test. You can choose to have a cervical screening test either by:

- taking your own sample from your vagina, using a simple swab (self-collection) or
- having a healthcare provider collect your sample from the cervix using a speculum.

Both options are covered by Medicare. If you decide to collect your own vaginal sample, you will be provided with a swab by your healthcare provider and instructions on how to collect the sample. You will be able to collect your sample in a private space. The sample is then sent to the laboratory where a genetic test is performed to detect the DNA of high-risk strains of HPV.

Which HPV strains are high risk?

- HPV types 6 and 11 typically cause venereal warts, but (along with types 42, 43, and 44) have a low risk of progressing to cancer.
- HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 68, 73 and 82 have a higher risk of progressing to cancer. HPV types 16 and 18 are associated with about 70 per cent of all cervical cancers.

If the test detects high-risk HPV virus, your sample will be sent to another department in the laboratory where a cytologist, a pathologist with special training in cervical cancer, will investigate it under the microscope to see whether your cervical cells are changing and becoming precancerous.



For more detailed information on these and many other tests go to pathologytestsexplained.org.au



www.pathologytestsexplained.org.au

Pathology Tests Explained is the primary national source of consumer information on pathology testing. Information is written and edited by practising pathologists and scientists, including leading experts. This ensures integrity and accuracy.

Pathology Tests Explained is managed by a consortium of medical and scientific organisations representing pathology practice in Australia. More details at:

www.pathologytestsexplained.org.au/about



My Health Record

You'll find a direct link to the Pathology Tests Explained website embedded in the pathology results pages of your My Health Record.

Click on the link to find information about what your tests are investigating or measuring and what your results can tell your doctor.

Please use this QR code to